

Original Article

Diagnostic value of Elastography in thyroid nodules

Akbar Aliasgarzadeh¹, Mohammad Kazem Tarzamani², Morteza Raeisi^{*3}, Majid Mobasseri¹, Maryam Mozayyan⁴, Morteza Ghojzadeh⁵

¹ Associate Professor, Bone Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

² Professor, Department of Radiology, School of Medicine, Imam Reza Hospital, Tabriz University of Medical Sciences, Tabriz, Iran

³ Medical Researcher, Bone Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁴ Medical Researcher, Department of Radiology, School of Medicine, Imam Reza Hospital, Tabriz University of Medical Sciences, Tabriz, Iran

⁵ Associate Professor, Liver and Gastrointestinal Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Article info

Article History:

Received: 22 Jan. 2014

Accepted: 27 Mar. 2014

Keywords:

Thyroid Nodule,
Ultrasonography,
Elastography

Abstract

Introduction: Palpable thyroid nodules are very common in the general population. Most of these nodules are benign and only about 5% of cases are malignant. Ultrasound elastography is a newly developed technique that uses ultrasound to provide an estimation of tissue stiffness. The aim of this study was to assess the diagnostic value of elastography in the evaluation of thyroid nodules supplementary to other ultrasonographic criteria.

Methods: In a descriptive cross-sectional study, thyroid nodules in 60 patients, who were candidates for fine needle aspiration biopsy (FNAB), were examined with conventional ultrasonography and elastography. The number and size of nodules, echogenicity, and presence of microcalcification, circumferential halo, and internal structure, margin of nodules, vascularity, and elasticity were determined. Pathologic evaluation of nodules was performed using FNAB. Imaging findings were compared with FNAB results.

Results: In the present study, 54 (90%) nodules were reported as soft and 6 (10%) as hard by ultrasound elastography. According to the results of FNAB, 9.5% were considered as non-benign nodules requiring surgery; all of these nodules were reported as soft in elastographic evaluation.

Conclusion: Routine elastography for all of the patients with thyroid nodule may not help the process of diagnosis. Further studies are recommended on its application in the management of thyroid nodules.

Citation: Aliasgarzadeh A, Tarzamani MK, Raeisi M, Mobasseri M, Mozayyan M, Ghojzadeh M. **Diagnostic value of Elastography in thyroid nodules.** *J Anal Res Clin Med* 2014; 2(2): 71-6.

Introduction

Palpable thyroid nodules are present in 4–7% of the population.¹⁻⁴ Most of these nodules are benign and only about 5% of cases are malignant.⁵⁻⁷

In patients who have thyroid nodule, fine needle aspiration biopsy (FNAB) is a useful tool for the diagnosis of thyroid cancer.⁸ It should be noted that FNAB is an invasive procedure and, in addition to the possibility of errors in sampling and analysis, this method is not cost effective for studying of all nodules.^{9,10}

Ultrasonography is the imaging study of choice for thyroid nodules. In addition to assessing the size and characteristics of the nodule, ultrasonography also provides guidance for diagnostic procedures such as FNAB. In comprehensive terms, ultrasound has a major role in studying thyroid nodules.³

Some sonographic features can be associated with increased risk of malignancy, including nodule hypoechogenicity, microcalcifications, central vascularity, irregular margins, an incomplete or absent

* Corresponding Author: Morteza Raeisi, E-mail: raiisy@yahoo.com

© 2014 The Authors; Tabriz University of Medical Sciences

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

halo, and an anteroposterior diameter greater than transverse diameter.^{11,12}

Ultrasound elastography is a newly developed technique that uses ultrasound to provide an estimation of tissue stiffness by measuring the degree of distortion under the application of an external force.¹³⁻¹⁵ There are several reports that this method can be helpful in distinguishing benign nodules from those that are malignant and in previous studies, the high stiffness of thyroid lesions has had correlation with malignancy.^{16,17}

Elastography has already been used to differentiate cancers from benign lesions in prostate, breast, pancreas, and lymph nodes.¹⁸⁻²¹ It has also been used for assessment of hepatic fibrosis.²² The aim of this study was to assess the diagnostic value of elastography in the evaluation of thyroid nodules supplementary to other ultrasonographic criteria.

Methods

This was a descriptive cross-sectional study. From the patients referred to the endocrinology outpatient clinics of the Tabriz University of Medical Sciences, Iran, from April to December 2013 with a diagnosis of thyroid nodule, 60 patients who were candidate for FNAB enrolled in this study. The inclusion criteria were the presence of palpable thyroid nodule and having normal thyroid function tests. The study group consisted of 52 women (86.7%) and 8 men (13.3%). The mean age of the patients was 45.83 ± 11.11 years (range: 22-76 years).

The study plan was reviewed and approved by the university ethical committee, and an informed consent was obtained from every patient. The project was funded by the Bone Research Center, Tabriz University of Medical Sciences.

Initially, B-mode and color Doppler ultrasound were performed for all nodules, by an experienced sonographer and then elastographic examination using Medison v10 ultrasound machine equipped with a 7.5 MHz linear probe was done for the prominent

nodules which were to undergo FNAB.

Conventional ultrasound images of thyroid nodules were obtained with patients positioned lying on their back with their neck slightly extended. The number and size of nodules, echogenicity, presence of microcalcification (tiny, punctuate echogenic foci of 1 mm or less), circumferential halo, internal structure, and margin of nodules were determined. Vascularity of each nodule was evaluated with color Doppler ultrasound.

The echogenicity of the nodules was described as hyperechoic, hypoechoic, and isoechoic in comparison to normal thyroid parenchyma. The nodules were categorized as solid, predominantly solid (< 50% cystic change), predominantly cystic (> 50% cystic change), and cystic based on their structure. The margins were classified as well-defined and poorly-defined. The shape was determined by measuring the anteroposterior diameter to transverse diameter ratio.

To perform elastography, the region of interest (ROI) was defined as a nodule with sufficient surrounding thyroid tissue. The technique used for the elastographic examination was the application of light compression with the ultrasound probe to the anterior neck.

As it is important to maintain a constant level of pressure throughout the examination and to reduce the intra-examiner variability, a standardized external compression was applied to the neck by using real time measurements displayed on an indicator-bar with a numerical scale (graded from 1 to 5) which gives a feedback on the compression quality.

In general, the nodules were classified into two groups according to the Ueno and Ito elasticity score; soft (elasticity in the whole or a large part of the nodule) and hard (elasticity only at the peripheral part of the nodule or no elasticity of the examined area).²³

Due to 7 patients' unwillingness to undergo FNAB, a total of 53 FNABs were performed.

If there were more than one nodule, FNAB was performed on that nodule which was

evaluated by elastography and had been identified by the sonographer.

FNAB was performed using a 23-gauge needle. Obtained specimens (6 to 10 slides), after fixation in alcohol, were sent for histopathologic examination.

To assess the interobserver agreement among pathologists in grading the quality of specimens, 10 specimens were analyzed by two different pathologists. Because of the excellent interobserver agreement between the two pathologists (kappa values > 0.7), one pathologist reported the rest of the specimens.

The Bethesda system was used for reporting thyroid cytopathology (Table 1).²⁴

The sonographer, FNAB performer, and pathologist were not aware of the purposes

of the study.

Statistical analysis was performed by SPSS for Windows (version 16; SPSS Inc., Chicago, IL, USA) using descriptive statistical methods.

Results

Among 60 patients, 29 (48.3%) patients had 1 thyroid nodule, 15 (25.0%) patients had 2, 6 (10.0%) patients had 3, and 10 (16.7%) patients had 4 or more than 4 thyroid nodules.

Sonographic findings of nodules are shown in table 2.

By ultrasound elastography, 54 (90%) nodules were reported as soft and 6 (10%) as hard. Results of pathologic evaluation are shown in table 1.

According to the results of FNAB,

Table 1. FNAB (fine needle aspiration biopsy) results, classified according to Bethesda system for reporting thyroid cytopathology

| Bethesda category | Frequency (n) | Percent (%) |
|---|---------------|-------------|
| I (nondiagnostic or unsatisfactory) | 6 | 11.3 |
| II (benign) | 41 | 77.4 |
| III (atypia of undetermined significance or follicular lesion of undetermined significance) | 1 | 1.9 |
| IV (follicular neoplasm or suspicious of a follicular neoplasm) | 2 | 3.8 |
| V (suspicions of malignancy) | 1 | 1.9 |
| VI (malignant) | 2 | 3.8 |
| Total | 53 | 100 |

Table 2. Sonographic findings of nodules

| Sonographic feature | Frequency (n) | Percent (%) | |
|------------------------------|------------------------------|-------------|------|
| Echogenicity | Hyperechoic | 31 | 51.7 |
| | Hypoechoic | 20 | 33.3 |
| | Isoechoic | 9 | 15.0 |
| Microcalcification | Present | 11 | 18.3 |
| | Absent | 49 | 81.7 |
| Margin | Sharp and well defined | 48 | 80.0 |
| | Irregular and poorly defined | 12 | 20.0 |
| Halo | Present | 15 | 25.0 |
| | Absent or incomplete | 45 | 75.0 |
| Texture (internal structure) | Solid | 21 | 35.0 |
| | Predominantly solid | 35 | 58.3 |
| | Predominantly cystic | 3 | 5.0 |
| Vascularity | Cystic | 1 | 1.7 |
| | Increased (central) | 13 | 21.7 |
| | Increased (peripheral) | 36 | 60.0 |
| Shape | Decreased | 10 | 16.7 |
| | Normal | 1 | 1.7 |
| Shape | Taller than wide | 7 | 11.7 |
| | Not taller than wide | 53 | 88.3 |

5 (9.5 %) of the 53 nodules were considered as non-benign nodules requiring surgery (Bethesda categories IV, V, and VI).

Sonographic features that imply an increased risk of malignancy in thyroid nodules, in 5 nodules which were considered as non-benign nodules requiring surgery, were as follows:

- 3 nodules were reported as hyperechoic, 1 as hypoechoic, and 1 isoechoic
- 4 nodules were without halo
- 2 nodules had a solid structure and 3 nodules were predominantly solid
- 3 nodules showed peripheral vascularity and 2 nodules had decreased vascularity
- Microcalcification was detected only in 1 nodule
- All 5 nodules had well-defined margins

It should be noted that all 5 of the above-mentioned nodules were reported as soft in elastographic evaluation.

Discussion

Thyroid nodules are relatively common and carry a low, but noticeable risk of malignancy.^{1,4,5} The most important challenge is differentiating benign from malignant nodules, and precise diagnosis and management of malignant nodules in the early stages.

The value of ultrasonography in the evaluation of thyroid nodules has been investigated in several studies.^{7,11,12}

In the latest guidelines released by the American Thyroid Association, if the thyroid function tests are normal, the next step in the work-up of thyroid nodule detected by palpation is ultrasonography.²⁵ Therefore, ultrasound is increasingly used for the evaluation of thyroid nodules.

It must be noted that the results of ultrasound are highly dependent on the instrument used and the operator. Specific sonographic criteria have been proposed to differentiate benign from malignant nodules.^{3,6,7} However, ultrasonography is not accepted in any reference as the sole method for making a final decision in approach to

thyroid nodule. Moreover, according to the American Thyroid Association recommendations, FNAB should ultimately be used as the procedure of choice for evaluating thyroid nodules.^{25,26}

Efforts are employed to avoid unnecessary FNABs by defining more precise criteria differentiating malignant from benign thyroid nodules with noninvasive methods. In this regard, elastography has been introduced as a technique that evaluates the degree of distortion of a tissue under the application of an external force. Because softer parts of tissue deform more readily than the stiffer parts, this technique enables objective evaluation of tissue stiffness from the deformation rate.²⁷ Furthermore, as noted, its performance has already been shown in the differential diagnosis of breast and prostate carcinomas.^{13,19}

Our study investigated the supplementary role of elastography in the differential diagnosis of thyroid nodules.

In the present study, 3.8% of thyroid nodules were malignant (Bethesda category VI) and 5.7% were suspicious for malignancy (Bethesda categories IV and V) based on the results of FNAB. This is in accordance with many studies.²⁴

In a retrospective evaluation of 849 nodules that were diagnosed at surgery or biopsy in 831 patients, the presence of at least 1 malignant feature had a sensitivity and specificity of 83.0% and 74.0%, respectively.²⁸ In another study of 1244 nodules in 900 patients, of 233 nodules that were ultrasonographically evaluated as malignant, 76.8% were also evaluated as malignant by cytology FNAB.²⁶

Elastographic findings in our study were not able to differentiate between benign and malignant nodules, but in other studies, including the study of Rago et al., elastography has shown great potential in the diagnosis of thyroid cancer, especially in nodules with indeterminate cytology.⁷

Asteria et al. have reported sensitivity of 94% and specificity of 81% for the differential diagnosis of thyroid nodules in elastographic

evaluation of 86 nodules in 67 patients.¹⁷ Kagoya et al., in a study performed on 47 thyroid nodules, have concluded that although elastography can assist in the differential diagnosis of thyroid nodules, its diagnostic performance is not ideal at present.²⁹

The noticeable point in some past studies is previous bias in sample selection. For example, samples have been selected from patients who were referred for surgery, but in our study, attempts have been made to enroll patients without any previous selection from the beginning of study, only if there is a nodule requiring initial evaluation.

Based on the results of our study, routine elastography may not help the process of diagnosis for all of the patients with thyroid nodule. If elastography is performed in selected cases, such as indeterminate or suspicious FNABs, it would probably be able to yield more applicable results.

The present study has some limitations. In our study, results of ultrasonography has little predictive value in differentiating benign and malignant thyroid nodules which seems to be due to the small sample size of the study, electing subjects from the entire referred population without any preliminary selection, and as a result, the low number of malignant nodules.

References

1. Hegedus L. Clinical practice. The thyroid nodule. *N Engl J Med* 2004; 351(17): 1764-71.
2. Cappelli C, Castellano M, Pirola I, Gandossi E, De ME, Cumetti D, et al. Thyroid nodule shape suggests malignancy. *Eur J Endocrinol* 2006; 155(1): 27-31.
3. Moon WJ, Baek JH, Jung SL, Kim DW, Kim EK, Kim JY, et al. Ultrasonography and the ultrasound-based management of thyroid nodules: consensus statement and recommendations. *Korean J Radiol* 2011; 12(1): 1-14.
4. Bomeli SR, LeBeau SO, Ferris RL. Evaluation of a thyroid nodule. *Otolaryngol Clin North Am* 2010; 43(2): 229-38, vii.
5. Yeung MJ, Serpell JW. Management of the solitary thyroid nodule. *Oncologist* 2008; 13(2): 105-12.
6. Hoang JK, Lee WK, Lee M, Johnson D, Farrell S. US Features of thyroid malignancy: pearls and pitfalls. *Radiographics* 2007; 27(3): 847-60.
7. Rago T, Santini F, Scutari M, Pinchera A, Vitti P. Elastography: new developments in ultrasound for predicting malignancy in thyroid nodules. *J Clin Endocrinol Metab* 2007; 92(8): 2917-22.
8. Screatton NJ, Berman LH, Grant JW. US-guided core-needle biopsy of the thyroid gland. *Radiology* 2003; 226(3): 827-32.
9. Hamburger JI. Diagnosis of thyroid nodules by fine needle biopsy: use and abuse. *J Clin Endocrinol Metab* 1994; 79(2): 335-9.
10. Castro MR, Gharib H. Continuing controversies in the management of thyroid nodules. *Ann Intern Med* 2005; 142(11): 926-31.
11. Cappelli C, Castellano M, Pirola I, Cumetti D, Agosti B, Gandossi E, et al. The predictive value of ultrasound findings in the management of thyroid nodules. *QJM* 2007; 100(1): 29-35.
12. Sipos JA. Advances in ultrasound for the diagnosis and management of thyroid cancer. *Thyroid* 2009; 19(12): 1363-72.

Conclusion

Ultrasonography and elastography may not replace cytologic evaluation in the assessment of thyroid nodule, and FNAB cytology is considered the gold standard diagnostic test for the diagnosis of thyroid nodules. Despite the value of elastography in differentiating benign from malignant lesions in other tissues debated in previous studies, in this study, elastography was not able to differentiate benign from malignant thyroid nodules.

Due to the noninvasive nature of ultrasound and elastography and easiness of performing them, further studies should be designed and implemented for their application in the management of thyroid nodules, considering more samples without sample selection bias. Conducting more surveys on elastography of thyroid nodules in suspicious cases prior to surgery will probably yield more tangible results.

Conflict of Interests

Authors have no conflict of interest.

Acknowledgements

The Authors would like to thank all the individuals and hospitals who participated in this study.

13. Itoh A, Ueno E, Tohno E, Kamma H, Takahashi H, Shiina T, et al. Breast disease: clinical application of US elastography for diagnosis. *Radiology* 2006; 239(2): 341-50.
14. Park SH, Kim SJ, Kim EK, Kim MJ, Son EJ, Kwak JY. Interobserver agreement in assessing the sonographic and elastographic features of malignant thyroid nodules. *AJR Am J Roentgenol* 2009; 193(5): W416-W423.
15. Dighe M, Bae U, Richardson ML, Dubinsky TJ, Minoshima S, Kim Y. Differential diagnosis of thyroid nodules with US elastography using carotid artery pulsation. *Radiology* 2008; 248(2): 662-9.
16. Lew JI, Solorzano CC. Use of ultrasound in the management of thyroid cancer. *Oncologist* 2010; 15(3): 253-8.
17. Asteria C, Giovanardi A, Pizzocaro A, Cozzaglio L, Morabito A, Somalvico F, et al. US-elastography in the differential diagnosis of benign and malignant thyroid nodules. *Thyroid* 2008; 18(5): 523-31.
18. Garra BS, Cespedes EI, Ophir J, Spratt SR, Zuurbier RA, Magnant CM, et al. Elastography of breast lesions: initial clinical results. *Radiology* 1997; 202(1): 79-86.
19. Cochlin DL, Ganatra RH, Griffiths DF. Elastography in the detection of prostatic cancer. *Clin Radiol* 2002; 57(11): 1014-20.
20. Lyshchik A, Higashi T, Asato R, Tanaka S, Ito J, Hiraoka M, et al. Cervical lymph node metastases: diagnosis at sonoelastography--initial experience. *Radiology* 2007; 243(1): 258-67.
21. Giovannini M, Hookey LC, Bories E, Pesenti C, Monges G, Delpero JR. Endoscopic ultrasound elastography: the first step towards virtual biopsy? Preliminary results in 49 patients. *Endoscopy* 2006; 38(4): 344-8.
22. Sandrin L, Fourquet B, Hasquenoph JM, Yon S, Fournier C, Mal F, et al. Transient elastography: a new noninvasive method for assessment of hepatic fibrosis. *Ultrasound Med Biol* 2003; 29(12): 1705-13.
23. Ueno E, Ito A. Diagnosis of breast cancer by elasticity imaging. *Eizo Joho Medical* 2004; 36: 2-6.
24. Cibas ES, Ali SZ. The Bethesda System for Reporting Thyroid Cytopathology. *Thyroid* 2009; 19(11): 1159-65.
25. Cooper DS, Doherty GM, Haugen BR, Kloos RT, Lee SL, Mandel SJ, et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2009; 19(11): 1167-214.
26. Ito Y, Amino N, Yokozawa T, Ota H, Ohshita M, Murata N, et al. Ultrasonographic evaluation of thyroid nodules in 900 patients: comparison among ultrasonographic, cytological, and histological findings. *Thyroid* 2007; 17(12): 1269-76.
27. Ophir J, Cespedes I, Ponnekanti H, Yazdi Y, Li X. Elastography: a quantitative method for imaging the elasticity of biological tissues. *Ultrason Imaging* 1991; 13(2): 111-34.
28. Moon WJ, Jung SL, Lee JH, Na DG, Baek JH, Lee YH, et al. Benign and malignant thyroid nodules: US differentiation--multicenter retrospective study. *Radiology* 2008; 247(3): 762-70.
29. Kagoya R, Monobe H, Tojima H. Utility of elastography for differential diagnosis of benign and malignant thyroid nodules. *Otolaryngol Head Neck Surg* 2010; 143(2): 230-4.